

UCLA

UCLA Previously Published Works

Title

Stereospecific Ring Contraction of Bromocycloheptenes through Dyotropic Rearrangements via Nonclassical Carbocation-Anion Pairs.

Permalink

<https://escholarship.org/uc/item/6hv54465>

Journal

Journal of the American Chemical Society, 140(15)

ISSN

0002-7863

Authors

Goh, Shermin S
Champagne, Pier Alexandre
Guduguntla, Sureshbabu
et al.

Publication Date

2018-04-01

DOI

10.1021/jacs.8b00821

Peer reviewed

Stereospecific Ring Contraction of Bromocycloheptenes through Dyotropic Rearrangements via Nonclassical Carbocation–Anion Pairs

Shermin S. Goh,^{†,||,#} Pier Alexandre Champagne,^{‡,#} Sureshbabu Guduguntla,[†] Takashi Kikuchi,^{§,⊥} Makoto Fujita,[§] K. N. Houk,^{*,‡,§} and Ben L. Feringa^{*,†,§}

[†]Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

[‡]Department of Chemistry and Biochemistry, University of California—Los Angeles, 607 Charles E. Young Drive East, Box 951569, Los Angeles, California 90095-1569, United States

[§]Department of Applied Chemistry, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan


[⊥]Rigaku Corporation, 3-9-12 Matsubara-cho, Akishima-shi, Tokyo 196-8628, Japan

Supporting Information

ABSTRACT: Experimental and theoretical evidence is reported for a rare type I dyotropic rearrangement involving a [1,2]-alkene shift, leading to the regio- and stereospecific ring contraction of bromocycloheptenes. This reaction occurs under mild conditions, with or without a Lewis acid catalyst. DFT calculations show that the reaction proceeds through a nonclassical carbocation–anion pair, which is crucial for the low activation barrier and enantiospecificity. The *chiral* cyclopropylcarbinyl cation may be a transition state or an intermediate, depending on the reaction conditions.

In the course of our investigation of the desymmetrization of *meso*-3,7-dibromocycloheptene **1**, we discovered that the initially formed enantiomerically enriched homoallylic bromides **2** were spontaneously isomerized with retention of configuration to form chiral substituted cyclohexenes **3** (Scheme 1).¹ This reaction could be accelerated by Lewis

Scheme 1. Desymmetrization of **1** by AAS and Subsequent Rearrangement of Cycloheptenes **2** to Cyclohexenes **3**^{a,b}



^aIsolated yields over two steps. ^b*er*'s of **2a–e** and **3a–e** were determined by chiral GC to be the same, so *es* is >99%.

acids such as silica or ZnBr₂, with no change in stereoselectivity. We have studied this reaction by a combination of experiments and computations, and now report that this involves a rare dyotropic rearrangement involving nonclassical cyclopropylcarbinyl cations on the reaction paths as either transition states or intermediates, depending on the conditions.

The copper-catalyzed desymmetrization of **1** by asymmetric allylic substitution² (AAS) with organolithium reagents³ initially afforded the expected products **2a–e** (>99:1 *dr*), as observed by NMR spectroscopy of the crude reaction mixtures [see Supporting Information (SI)]. The reaction also proceeded with high enantioselectivity, as determined by chiral GC.¹ However, upon exposure to silica, the bromocycloheptenes **2a–e** isomerized to afford six-membered cyclic homoallylic bromides **3a–e**. The ring contraction reaction proceeded with complete regioselectivity and enantiospecificity, as determined by chiral GC [see SI, part 1 and SI, part 2].

In fact, the rearrangement reaction was so facile that the transformation from cycloheptenes **2a–e** to cyclohexenes **3a–e** occurred even on neutral alumina, or when left standing in chloroform-*d* (*t*_{1/2} ≈ 6 days). The major enantiomer of **3d**,⁴ obtained as a colorless oil, was determined to be (*R,R*) by X-ray crystallography using only 5 μg of compound and the crystalline sponge method developed by the Fujita group⁵ (Figure 1). A similar attempt was made to study the absolute




Figure 1. Ball-and-stick representation of the X-ray structure of **3d** determined by the crystalline sponge method.

configuration of its precursor **2d**; however, the rearrangement occurred while soaking in crystalline sponge, and only **3d** was detected when the X-ray analysis was performed.

The ring contraction reaction from **2a–e** to **3a–e**, interconverting the two isomeric homoallylic bromides, involves the 1,2-positional exchange of the alkenyl and bromo groups; thus it is formally a [2,2]-dyotropic rearrangement. These were first described in 1972 by M. T. Reetz as a class of pericyclic valence isomerizations involving the simultaneous


Received: January 23, 2018

Published: March 29, 2018


intramolecular migration of two σ -bonds.^{6,7} Type I dyotropic rearrangements occur when the two migrating groups exchange their positions (e.g., the classic rearrangement of *anti* vicinal dibromides, Scheme 2a),⁸ while in type II rearrangements the

Scheme 2. Uncatalyzed Concerted Type I Dyotropic Rearrangements Occurring on a Static C–C Scaffold

(a) Vicinal dibromide dyotropic rearrangement (Ref 8)



(b) Alkyl-mesylate dyotropic rearrangement (Ref 20)



groups migrate to entirely different positions. Dyotropic reactions have become more common recently in organic and organometallic chemistry⁹ and have even been applied to total syntheses.^{9,10} The concerted migration of carbon chains through a [2,2]-shift on a static C–C scaffold is uncommon. [σ_2 + σ_2] processes are thermally forbidden by the Woodward–Hoffmann rules,^{11,12} but the reaction becomes allowed if one migrating group has a lone pair and migrates with inversion. Nevertheless, such reactions are still rare because the activation barriers are usually quite high.¹³

Until recently, the only experimental examples of such reactions involved highly strained lactones reacting under strenuous conditions. Examples of alkyl group migration are the ring expansion of β -lactones to butyrolactones promoted by stoichiometric magnesium bromide,^{14–16} or the rearrangement of cage δ -lactones to γ -lactones at 350 °C on a quartz column.¹⁷ Acyl group migration involving a β -lactone ring expansion

promoted by stoichiometric Lewis acid is also known.¹⁸ In recent years, milder dyotropic rearrangements have been discovered. Gutta and Tantillo proposed a 1,2-positional exchange of an alkyl group and a hydrogen atom in their computed biosynthetic pathway for formation of pentalenene in 2006.¹⁹ More recently, Faza, Lopez, and co-workers reported the type I dyotropic ring expansion of hydrindane to decalin occurring at –78 °C upon mesylation (Scheme 2b).²⁰ To the best of our knowledge, a concerted and uncatalyzed type I dyotropic migration of a C–C π system is still unknown.

The mechanism of this rearrangement was investigated by NMR spectroscopy. Time-dependent aliquot studies with silica as reagent (500% w/w) for the rearrangement of bromocycloheptene **2d** showed that bromocyclohexene **3d** (>99% *es*) was the only product generated. This was observed in both polar aprotic (chloroform-*d*) and apolar (benzene-*d*₆) solvents, although the rate of reaction was slower in benzene-*d*₆ (Figure 2b,c); the lack of solvent dependence for selectivity indicates that the reaction does not proceed through a discrete carbocation. The use of Lewis acids (e.g., ZnBr₂ and TMSOTf) led to poorer selectivity in the rearrangement reaction, with ca. 5–10% of a different diastereoisomer **4d** observed (Figure 2d,e);²¹ its formation may be explained by the ionization of product **3d** by the stronger Lewis acids [see SI]. There was, fortunately, no erosion in *ee* of the expected product. These reactions with Lewis acids could be monitored *in situ* by NMR time course experiments in either chloroform-*d* (Figure 2a) or benzene-*d*₆ with no difference in product ratio or enantiospecificity, although the reaction was always slower in the apolar solvent [see SI]. The reaction could also be performed with Brønsted acids [see SI], or with catalytic amounts of Lewis acids (Figure 2e), albeit at a much reduced rate. When conducted in the presence of radical scavenger BHT (1.0 equiv), the reaction profile did not change [see SI], indicating that a radical pathway was not involved.

To further understand the mechanism and specificity of the rearrangement, we performed DFT calculations on compound




Figure 2. Left: (a) ¹H NMR spectroscopic profiles of bromocycloheptene **2d** with ZnBr₂ (1 equiv) in chloroform-*d* with increasing time; reaction scheme is shown at top. Right: Reaction progress of **2d** with silica (500% w/w) monitored by timed aliquots in (b) chloroform-*d* and (c) benzene-*d*₆. Reaction progress of **2d** monitored by *in situ* NMR spectroscopy with (d) ZnBr₂ (1 equiv) in chloroform-*d* and (e) TMSOTf (0.5 equiv) in chloroform-*d*.

2a at the M06-2X/6-311+G(d,p)//M06-2X/6-31G(d) level of theory,²² using Gaussian 09.²³ The SMD solvation model²⁴ for CHCl₃ was used throughout. Computed structures were visualized using CYLview.²⁵ We first investigated whether a concerted uncatalyzed dyotropic rearrangement is plausible. The cycloheptene ring of **2a** has two low-energy conformations, where the methyl and bromo substituents are either pseudoaxial (**2a-ax**) or pseudoequatorial (**2a-eq**). From **2a-eq**, a dyotropic rearrangement involves the experimentally observed 1,2-shift of both the alkenyl and bromo groups to form **3a**. The barrier for this transformation is only 25.2 kcal/mol (Figure 3), and the reaction is exergonic by 5.0 kcal/mol.




Figure 3. Calculated free energy profile for the concerted dyotropic rearrangements from the pseudoaxial (left part) or pseudoequatorial (right part) conformations of **2a**.

In contrast, the dyotropic rearrangement of the methyl and bromo groups from **2a-ax**, forming **5a**, has a significantly higher barrier of 40.1 kcal/mol. The calculated ΔG^\ddagger for TS-eq is consistent with our observations of a slow uncatalyzed transformation of **2a-e** to **3a-e** at room temperature in chloroform ($t_{1/2} \approx 6$ days).

Both TS-eq and TS-ax are highly polarized, with two long C–Br bonds (>3.0 Å) and two short C–C bonds, and as such can be described as tight ion-pairs of a carbocation and bromide anion. Indeed, the bromine atom in these two TSs bears an almost full negative charge (−0.95 to −0.97 au, see SI). This is in stark contrast to classical type I dyotropic rearrangements of vicinal dibromides, where the C–Br bonds were calculated to be around 2.5–2.8 Å.^{8f} It is known that delocalization of the formed π bond in the TSs of dyotropic rearrangements provides key stabilization.^{8d,e} In the case of TS-eq, the large polarization causes the carbon backbone of the substrate to approach the geometry of a $\pi\sigma$ -delocalized bisected cyclopropylcarbinyll (nonclassical) cation (vide infra).^{26,27} This greatly stabilizes the π bond of the TS (in blue, Figure 3) and explains why TS-eq has such a low barrier compared to TS-ax, for which the carbon backbone has the geometry of a less-stabilized corner-protonated cyclopropane.²⁸

Having established that the enantiospecific uncatalyzed reaction is very likely to operate through a concerted dyotropic rearrangement, we investigated the role of an external Lewis acid (Figure 4). Formation of **2a·ZnBr₂** from the isolated reactants is favorable by 8.9 kcal/mol [see SI]. From this coordinated species, no concerted dyotropic transition states could be located; instead a stepwise mechanism is found.




Figure 4. Calculated free energy profile for the stepwise transformation of **2a** to **3a**, catalyzed by ZnBr₂. For the structure of **6a·ZnBr₃**, the ZnBr₃[−] is hidden for clarity.

Ionization of the C–Br bond of complex **2a·ZnBr₂** through TS 1 has a low barrier of 14.6 kcal/mol and leads to a contact ion-pair consisting of nonclassical cyclopropylcarbinyll cation **6a** and ZnBr₃[−]. From this intermediate, only 7.8 kcal/mol is required to reach TS 2, where the C–Br bond of **3a** is formed. These calculations are consistent with the much faster reaction observed when ZnBr₂ is used as Lewis acid, since the activation barrier is predicted to be almost 10 kcal/mol lower than in the uncatalyzed case (14.6 vs 25.2 kcal/mol). While some stepwise ionic dyotropic rearrangements are known, these were limited to intramolecular examples (e.g., dyotropic rearrangement of Himbert cycloadducts via a zwitterion).^{29,30} The present mechanism is an unprecedented case of a stepwise formal dyotropic rearrangement proceeding through an ion-pair.

Ionization of the C–Br bond in TS 1 happens with simultaneous backside attack of the alkenyl group, analogous to an S_N2 mechanism. Similarly, bromide attack in TS 2 occurs with release of the alkenyl group; therefore the C–Br bonds are broken and formed from the same face of **6a**, the structure of which is determined by the stereochemistry of **2a**. Thus **6a** is a *chiral* carbocation; however, even in the presence of Lewis acids such as ZnBr₂, the reaction of **2d** to **3d** was shown to be perfectly enantiospecific. We performed additional calculations to investigate the possible mechanism of racemization of **6a**. It is now well established that cyclopropylcarbinyll cations^{26,27} are in equilibrium with the related bicyclobutonium cations, with the latter being more stable for the parent C₄H₇⁺.^{31,32} For the bicyclic cyclopropylcarbinyll cation **6a**, the profile is much more complex. The lowest-energy path for the racemization of **6a** is through the *meso*-bicyclobutonium **7a**, which is a TS (instead of a minimum) on the potential energy surface of cation **6a** (Figure S).³³

Once *chiral* cation **6a** is formed from the C–Br bond cleavage of **2a**, it requires an additional 10.9 kcal/mol of free energy to racemize through **7a·ZnBr₃**, a bicyclobutonium ion-pair. In contrast, it only needs 7.8 kcal/mol to recombine with the ZnBr₃[−] counteranion to form the stable product **3a** through TS 2. Moreover, the latter reaction is not reversible, as **3a·ZnBr₂** is 5.4 kcal/mol more stable than the starting complex. As such, there is a kinetic barrier to racemization in this system, allowing a stepwise enantiospecific rearrangement to occur.




Figure 5. Lowest-energy pathway for racemization of **6a•ZnBr₃**. Free energies (kcal/mol) are relative to **2a•ZnBr₂** (Figure 4). The ZnBr_3^- anions are hidden for clarity.

In summary, we have discovered an unanticipated ring contraction of bromocycloheptenes under mild conditions with remarkable regio- and stereochemistry. The ring contraction occurs via a double 1,2-migration of an alkene group and a bromide. DFT calculations show that the reaction proceeds through a nonclassical carbocation–anion pair; the $\pi\sigma$ -delocalized bisected cyclopropylcarbanyl cation is crucial for the low activation barrier and enantiospecificity of the rearrangement. The reaction is concerted when uncatalyzed, presenting a rare type I dyotropic rearrangement of an alkene on a C–C stationary scaffold. In contrast, the reaction follows a stepwise mechanism under Lewis acid catalysis, and can be described as a formal dyotropic rearrangement. Our study also highlights how cations derived from chiral homoallylic halides may be productive intermediates in other enantiospecific reactions.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.8b00821.

Experimental procedures and characterization data, and data for computed structures (PDF)

NMR and chiral GC spectra (PDF)

X-ray crystallographic data for compound **3d** (CIF)

■ AUTHOR INFORMATION

Corresponding Authors

*houk@chem.ucla.edu

*b.l.feringa@rug.nl

ORCID

Pier Alexandre Champagne: 0000-0002-0546-7537

Makoto Fujita: 0000-0001-6105-7340

K. N. Houk: 0000-0002-8387-5261

Ben L. Feringa: 0000-0003-0588-8435

Present Address

^{||}S.S.G.: Institute of Materials Research and Engineering, 2 Fusionopolis Way, Innovis #08-03, Singapore 138634

Author Contributions

[#]S.S.G. and P.A.C. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

B.L.F. acknowledges The Netherlands Organization for Scientific Research (NWO-CW), the Royal Netherlands Academy of Arts and Sciences (KNAW), and the Ministry of Education Culture and Science (Gravitation program 024.601035) for funding. S.S.G. acknowledges A*STAR (NSS) for a postdoctoral fellowship. P.A.C. acknowledges the FRQNT for a postdoctoral fellowship. We are grateful to the NSF (CHE-1361104 to K.N.H.) for financial support of this research. Computations were performed on the Hoffman2 cluster at UCLA. M.F. and T.K. acknowledge JST-ACCEL project in which M.F. is a principal investigator.

■ REFERENCES

- (1) Desymmetrization of *meso*-dibromocycloalkenes with organolithium reagents: unpublished manuscript.
- (2) For reviews of Cu-catalyzed AAS, see: (a) Geurts, K.; Fletcher, S. P.; van Zijl, A. W.; Minnaard, A. J.; Feringa, B. L. *Pure Appl. Chem.* **2008**, *80*, 1025–1037. (b) Harutyunyan, S. R.; den Hartog, T.; Geurts, K.; Minnaard, A. J.; Feringa, B. L. *Chem. Rev.* **2008**, *108*, 2824–2852. (c) Alexakis, A.; Bäckvall, J. E.; Krause, N.; Pàmies, O.; Diéguez, M. *Chem. Rev.* **2008**, *108*, 2796. (d) Langlois, J.-B.; Alexakis, A. In *Transition Metal Catalyzed Allylic Substitution in Organic Synthesis*; Kazmaier, U., Ed.; Springer-Verlag: Berlin, 2012; pp 235–268. (e) Hornillos, V.; Gualtierotti, J.-B.; Feringa, B. L. *Top. Organomet. Chem.* **2016**, *58*, 1–39. (f) Baslé, O.; Denicourt-Nowicki, A.; Crévisy, C.; Mauduit, M. In *Copper-Catalyzed Asymmetric Synthesis*; Alexakis, A.; Krause, N.; Woodward, S., Eds.; Wiley-VCH: Weinheim, 2014; pp 85–126.
- (3) For selected examples of asymmetric allylic substitution with organolithium reagents, see: (a) Pérez, M.; Fañanás-Mastral, M.; Bos, P. H.; Rudolph, A.; Harutyunyan, S. R.; Feringa, B. L. *Nat. Chem.* **2011**, *3*, 377. (b) Fañanás-Mastral, M.; Pérez, M.; Bos, P. H.; Rudolph, A.; Harutyunyan, S. R.; Feringa, B. L. *Angew. Chem., Int. Ed.* **2012**, *51*, 1922. (c) Guduguntla, S.; Gualtierotti, J.-B.; Goh, S. S.; Feringa, B. L. *ACS Catal.* **2016**, *6*, 6591. (d) Hornillos, V.; Guduguntla, S.; Fañanás-Mastral, M.; Pérez, M.; Bos, P. H.; Rudolph, A.; Harutyunyan, S. R.; Feringa, B. L. *Nat. Protoc.* **2017**, *12*, 493.
- (4) CCDC 1571256 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- (5) For selected examples of crystalline sponge X-ray crystallography, see: (a) Inokuma, Y.; Yoshioka, S.; Ariyoshi, J.; Arai, T.; Hitora, Y.; Takada, K.; Matsunaga, S.; Rissanen, K.; Fujita, M. *Nature* **2013**, *495*, 461. (b) Inokuma, Y.; Yoshioka, S.; Ariyoshi, J.; Arai, T.; Fujita, M. *Nat. Protoc.* **2014**, *9*, 246. (c) Hoshino, M.; Khutia, A.; Xing, H.; Inokuma, Y.; Fujita, M. *IUCr* **2016**, *3*, 139.
- (6) Reetz, M. T. *Angew. Chem., Int. Ed. Engl.* **1972**, *11* (2), 129.
- (7) Reetz, M. T. *Angew. Chem., Int. Ed. Engl.* **1972**, *11* (2), 130.
- (8) For selected experimental and theoretical studies on the dyotropic rearrangement of *anti* vicinal dibromides, see: (a) Grob, C. A.; Winstein, S. *Helv. Chim. Acta* **1952**, *35* (3), 782. (b) Barton, D. H. R.; Head, A. J. *J. Chem. Soc.* **1956**, 932. (c) Barili, P. L.; Bellucci, G.; Berti, G.; Marioni, F.; Marsili, A.; Morelli, I. *J. Chem. Soc. D* **1970**, 1437. (d) Frontera, A.; Suner, G. A.; Deya, P. M. *J. Org. Chem.* **1992**, *57* (25), 6731. (e) Zou, J.-W.; Yu, C.-H. *J. Phys. Chem. A* **2004**, *108*, 5649. (f) Fernández, I.; Sierra, M. A.; Cossio, F. P. *Chem. - Eur. J.* **2006**, *12* (24), 6323.
- (9) Fernández, I.; Cossio, F. P.; Sierra, M. A. *Chem. Rev.* **2009**, *109* (12), 6687.
- (10) For selected examples of total syntheses involving dyotropic rearrangements, see: (a) Kocienski, P.; Barber, C. *Pure Appl. Chem.* **1990**, *62* (10), 1933. (b) Lin, S.; Danishefsky, S. J. *Angew. Chem., Int. Ed.* **2002**, *41* (3), 512. (c) Li, W.; LaCour, T. G.; Fuchs, P. L. *J. Am.*

- Chem. Soc.* **2002**, 124, 4548. (d) Denmark, S. E.; Montgomery, J. I. *Angew. Chem., Int. Ed.* **2005**, 44 (24), 3732.
- (11) Reetz, M. T. *Tetrahedron* **1973**, 29 (14), 2189.
- (12) Reetz, M. T. *Adv. Organomet. Chem.* **1977**, 16, 33.
- (13) Gutierrez, O.; Tantillo, D. J. *J. Org. Chem.* **2012**, 77 (20), 8845.
- (14) Mulzer, J.; Brüttrup, G. *Angew. Chem., Int. Ed. Engl.* **1979**, 18 (10), 793.
- (15) Black, T. H.; DuBay, W. J.; Tully, P. S. *J. Org. Chem.* **1988**, 53 (25), 5922.
- (16) Arrastia, I.; Lecea, B.; Cossio, F. P. *Tetrahedron Lett.* **1996**, 37 (2), 245.
- (17) Satō, K.; Yamashita, Y.; Mukai, T. *Tetrahedron Lett.* **1981**, 22 (52), 5303.
- (18) Leverett, C. A.; Purohit, V. C.; Johnson, A. G.; Davis, R. L.; Tantillo, D. J.; Romo, D. *J. Am. Chem. Soc.* **2012**, 134 (32), 13348.
- (19) Gutta, P.; Tantillo, D. J. *J. Am. Chem. Soc.* **2006**, 128 (18), 6172.
- (20) Santalla, H.; Faza, O. N.; Gómez, G.; Fall, Y.; Silva López, C. *Org. Lett.* **2017**, 19 (13), 3648.
- (21) More experimental data for the rearrangement of **2d** to **3d** monitored by NMR spectroscopy can be found in the SI.
- (22) Zhao, Y.; Truhlar, D. G. *Theor. Chem. Acc.* **2008**, 120, 215.
- (23) Frisch, M. J.; et al. *Gaussian 09*, Revision D.01; Gaussian, Inc.: Wallingford, CT, 2013.
- (24) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. *J. Phys. Chem. B* **2009**, 113 (18), 6378.
- (25) Legault, C. Y. *CYLVview*, 1.0b; Université de Sherbrooke, 2009; <http://www.cylvview.org>.
- (26) Roberts, J. D.; Mazur, R. H. *J. Am. Chem. Soc.* **1951**, 73 (7), 3542.
- (27) Roberts, J. D.; Mazur, R. H. *J. Am. Chem. Soc.* **1951**, 73 (6), 2509.
- (28) (a) Dewar, M. J. S.; Healy, E. F.; Ruiz, J. M. *J. Chem. Soc., Chem. Commun.* **1987**, 943. (b) Burritt, A.; Coxon, J. M.; Steel, P. J. *J. Org. Chem.* **1995**, 60 (23), 7670.
- (29) Pham, H. V.; Karns, A. S.; Vanderwal, C. D.; Houk, K. N. *J. Am. Chem. Soc.* **2015**, 137 (21), 6956.
- (30) Davis, R. L.; Leverett, C. A.; Romo, D.; Tantillo, D. J. *J. Org. Chem.* **2011**, 76 (17), 7167.
- (31) Staral, J. S.; Yavari, I.; Roberts, J. D.; Prakash, G. K. S.; Donovan, D. J.; Olah, G. A. *J. Am. Chem. Soc.* **1978**, 100 (25), 8016.
- (32) Olah, G. A.; Surya Prakash, G. K.; Rasul, G. *J. Am. Chem. Soc.* **2008**, 130 (28), 9168.
- (33) The full free energy profile of **6a** can be found in the SI.